

A Computational Algorithm for $\pi(N)$

Abhijit Sen
Theory Group, Saha Institute of Nuclear Physics
1/AF, BidhanNagar, Calcutta-700064
West Bengal, INDIA

Satyabrata Adhikari
Calcutta Mathematical Society, A.E 374
BidhanNagar, Sector-I, Calcutta-700064
West Bengal, INDIA

Abstract

An algorithm for computing $\pi(N)$ is presented. It is shown that using a symmetry of odd composites we can easily compute $\pi(N)$. This method relies on the fact that counting the number of odd composites not exceeding N suffices to calculate $\pi(N)$.

Key Words: Prime Number, Odd Composite, Cardinality
AMS classification number(1980):10-01

Introduction

The problem of finding a compact mathematical formula for $\pi(N)$ is well known [1]. Though no compact result has been reported till date, a number of accurate/fairly accurate algorithms exist [2,3,7,8] among which the prime number theorem [4] and the function $\text{Li}(x)$ [3] require special mention. Wilson's theorem [5] seems to be unique, however, like all reportings so far it has a low practical utility. Recently a result has been reported using Smarandache Functions [6] which too however has a low practical utility.

In the present paper we develop a computational algorithm based on a certain symmetry of odd composites. This is the main essence of the present piece of work.

The Algorithm

A. General Formulation

Among all natural numbers we know that

- i) All even numbers except 2 are composite.
- ii) All odd composites can be factorised into odd primes only.
- iii) Unity is neither prime nor composite.

Therefore we have

$$\pi(N) = N - n_e(c) - n_o(c) - 1 \quad \dots\dots\dots (1)$$

where

$n_e(c)$ and $n_o(c)$ give respectively the number of even and odd composite numbers not exceeding N . Without the loss of generality we assume $N \geq 2$.

Now we have

$$n_e(c) = \text{Floor} \left(\frac{N}{2} - 1 \right) \quad \dots\dots\dots (1a)$$

To obtain $n_o(c)$ let us refer to the multiplication table of odd natural numbers ≥ 3 as shown in Table-II. It shows a very small section at the top left hand corner of the infinite multiplication table.

It can be shown that all odd composites occur at least once within Table - II whose general term is given by

$$t_{n,r} = (2n+1)^2 + 2(r-1)(2n+1) = (2n+r)^2 - (r-1)^2 \quad \dots\dots\dots (2)$$

where both n and r are natural.

We further see that for each 'n' value we have an A.P originating at $(2n+1)^2$ with a common difference $2(2n+1)$. The terms of the A.P for each 'n' are

assumed to constitute a set represented as

$$\{ t_{n,r} \} = \{ x : x = (2n + r)^2 - (r - 1)^2 \}$$

where r varies from 1 to

$$r|_{max} = \text{Floor} \left(\frac{1}{2(2n+1)} (N - (2n+1)^2) + 1 \right) \dots\dots\dots (3a)$$

for a given n i.e for a given set $\{ t_{n,r} \}$.

Thus,

$$n_o(c) = \sum \text{Floor} \left(\frac{N-(2n+1)^2}{2(2n+1)} + 1 \right) - \lambda_c \dots\dots\dots (4a)$$

where the sum is over n from 1 to

$$n|_{max} = \text{Floor} \left(\frac{\sqrt{N}-1}{2} \right) \dots\dots\dots (3b).$$

Here λ_c is a correction term which we would discuss shortly.

B. Corrections

B1. The Summation

For a composite $(2n_r+1)$, all the terms arising from the summation in (4a) corresponding to $n=n_r$ would be a duplication of some previously occurring term for some $n = n_s < n_r$. Therefore we shall restrict ourselves to only those values of n_r for which $(2n_r + 1)$ is prime.

Following the discussions as above we can recast the summation part of (4a) as

$$n_o(c) = \sum_i^R \left(\text{Floor} \frac{N-(2n_i+1)^2}{2*(2n_i+1)} + 1 \right) - \lambda_c \dots\dots (4b)$$

where we restrict the sum for a prime $(2n_i+1)$. Here the suffix R denotes the sum over the restricted sets of the values of n's. The primality of $(2n_i+1)$ can be ascertained using the symmetry we mentioned. We do not accept those values of n_k for which

$$\frac{1}{2} (t_{n,r} - 1) = n_k \leq n_{max} \text{ for } r \leq r_{max} \text{ as given by (3a) and (3b).}$$

As will be shown in a latter article [9] all composite $(2n_i + 1)$ can be expressed as a difference of two squares following eqn. (2) .

Alternatively the primality could be determined using Wilson's theorem[5].

B2. Evaluation of λ_c

The need of λ_c arises from the fact that for different n values the sets $\{ t_{n,r} \}$ coincide at certain points.

Now once the summation in (4a) be restricted we shall also require to restrict the evaluation of λ_c .

Restricting ourselves to those values of n_r for which $(2n_r+1)$ is prime we can evaluate λ_c using the set theoretic formula

$$n(A_1 \cup A_2 \cup \dots \cup A_k) = \sum n(A_i) - \sum \sum n(A_i \cap A_j) + \sum n(A_i \cap A_j \cap A_k) - \dots \quad (5)$$

where $n(X)$ denotes the cardinality of set X .

Here summations over all indices can range from 1 to n_{max} with the restriction $i < j < k \dots$ and so on.

Let us define the following notations:

For intersection of two A.P's let

$t_{n_1, n_2}^{(1)}$ be the first common term of $\{t_{n_1, r}\}$ and $\{t_{n_2, r}\}$ while

$(c.d)_{n_1, n_2}$ be the common difference of the common terms

For intersection of more than two A.P's the notation is identical.

It can be shown that

a) For two A.P's

$$t_{n_1, n_2}^{(1)} = (2n_1 + 1) * (2n_2 + 1) * \{3 + 2 * \text{Ceil}[Z] * H[Z]\} \dots \quad (6a)$$

where

$$Z = \frac{1}{2} \left(\frac{2n_2 + 1}{2n_1 + 1} - 3 \right)$$

and

$$(c.d)_{n_1, n_2} = 2 * (2n_1 + 1) * (2n_2 + 1) \dots \quad (6b)$$

Here $H[Z]$ denotes the Heaviside Step Function defined by

$$\begin{aligned} H[Z] &= 1, z > 0 \\ &= \frac{1}{2}, z = 0 \\ &= 0, z < 0 \end{aligned}$$

b) For more than two A.P's

$$t_{n_1, n_2 \dots n_k}^{(1)} = \prod_{i=1}^k (2n_i + 1) * (1 + 2 * \text{Ceil}(Q) * H[Q]) \dots \quad (7a)$$

where

$$Q = \frac{1}{2} \left(\frac{(2n_k + 1)^2}{\prod_{i=1}^k (2n_i + 1)} - 1 \right)$$

and

$$(c.d)_{n_1, n_2, \dots, n_k} = 2 * \prod_{i=1}^k (2n_i + 1) \dots \quad (7b)$$

Using (5), (6a), (6b), (7a) and (7b) we get

$$\lambda_c = \sum_{r=2}^{n_{max}} (-1)^r \sum^R \sum^R \dots \sum^R (\text{Floor}(B) + 1) * H[B + \varepsilon] \dots \quad (8)$$

where

i) The summations run over all restricted sets of n values, one for each set with

$n_1 < n_2 < n_3 \dots < n_k$

ii) ε is a positive number, however small. It will be neglected at the last stages.

iii) We have

$$B = \frac{N - t_{n_1, n_2, \dots, n_r}^{(1)}}{(c.d)_{n_1, n_2, \dots, n_r}}$$

The Heaviside function restricts the numerator to non-negative values.

Using (1),(1a),(4b),(5), and (8) the result follows.

Though equation (8) seems to have many steps in practice it would be quite limited as once the result given by (6a) and (7a) exceeds N, higher n_r terms would be in general very very limited in number.

The Results

The results as obtained from the above mentioned algorithm are given in Table I. It is seen that the value of $\pi(N)$ is exactly reproduced. This establishes the correctness of the above mentioned algorithm.

To illustrate the method let us evaluate $\pi(100)$. We have

from (3b) : $n_{max}=4$

Now, to calculate $\pi(100)$ we need to know all primes till $\sqrt{100}=10$.

Corresponding to $n=4$ we get $(2n+1)=9 < 10$ which is a composite.

So we do not consider $n=4$.

Proceeding;

from (1a) : $n_e(c) = \text{Floor}(\frac{100}{2} - 1) = 49$

from (4b) : $n_o(c) = 16 + 8 + 4 = 28$

from (8) : $\lambda_c = 2 + 1 = 3$

Therefore from (1) we have $\pi(100) = 100 - 49 - 28 + 3 - 1 = 25$

Conclusion

The formula for $\pi(N)$ as established seems complicated at first sight. However the symmetry of natural numbers used, the necessity of introduction of λ_c and the simple set-theoretic prescription for its evaluation makes the present method quite tractable. However, like the well known "Sieve algorithm" [8] this method too requires the knowledge of all primes till $\sqrt{N}$ to evaluate $\pi(N)$. The number of steps is much reduced as we use the symmetry of natural numbers throughout, first to determine primality of $(2n_i+1)$ if the same is not already known and afterwards for the final evaluation of $\pi(N)$ in comparison to the Sieve Algorithm. Referring back to the symmetry of odd composites that we had mentioned, the symmetry is more clearly evident if we use the binary base of representation specially in Table II. This is evident from the fact that the common difference of the common differences in Table II equals base squared. It is believed that this could be a suitable method to determine $\pi(N)$.

Acknowledgement

The authors would like to acknowledge the constant encouragement by Mr. Samit De of SINP.

TABLE : I

Results of determination of $\pi(N)$

N	$\pi(N)$	Estimated value
10	4	4
100	25	25
1000	168	168

TABLE : II

Multiplication Table of odd natural numbers

	3	5	7	9	11	Common Difference
3	9	15	21	27	33	6
5		25	35	45	55	10
7			49	63	77	14
Common Difference						$4 = 2^2$

References

1. A-17, Unsolved problems in Number Theory, Richard K Guy, Springer-Verlag, N.Y
2. CRC Concise Encyclopedia of Mathematics, Eric W Weisstein, 1630, Chapman & Hall/CRC.
3. CRC Concise Encyclopedia of Mathematics, Eric W Weisstein, 1097, Chapman & Hall/CRC.
4. Elementary Number Theory, David M. Burton, Appendixes, 396-403 (1995)
5. Math. Gaz. 48 (1964) 413-415
6. CRC Concise Encyclopedia of Mathematics, Eric W Weisstein, 1660 - 1661, Chapman & Hall/CRC.
7. CRC Concise Encyclopedia of Mathematics, Eric W Weisstein, 1427 - 1428, Chapman & Hall/CRC.
8. Elementary Number Theory, David M. Burton, 60-61 (1995)
9. <http://www.arXiv.org>, math.GM/0202269